# Five dimensional SUSY gauge theories in the context of M theory

Author:

Mohammed Hakeem

Supervisor:

Prof. Amihay Hanany



Submitted in part fulfilment of the requirements for the degree of Master of Science in Theoretical Physics of Imperial College London

September 21, 2014

# Acknowledgement

I would Like to thank Prof. Hanany for his distinguished introduction to string theory through his course, and for giving me the opportunity to investigate and review the interesting phenomenon that arise in the field of supersymmetric field theories and brane dynamics.

I would also like to thank my parents for their continued love and support, which is the driving force behind my academic achievements.

## Contents

.

1	Intr	oduction	3		
<b>2</b>	Fiel	d Theory content	5		
3	Type IIB setup				
	3.1	Vertices	8		
	3.2	Webs	9		
	3.3	Gauge theories	10		
	3.4	Deformations	17		
	3.5	Grid diagrams	19		
4	Curves 22				
	4.1	M theory	22		
	4.2	Complex coordinates	23		
	4.3	Curves from grids	26		
	4.4	General $SU(N_c)$	27		
	4.5	Four dimensions	30		
<b>5</b>	Con	clusion	32		

## 1 Introduction

The intertwining relationship between brane dynamics and non-Abelian gauge theories has long been an intuitive approach to understanding supersymmetric gauge theories in several dimensions. A substantial amount of work has been done in the past decade in realizing supersymmetric field theories, in [9] three dimensional theories were formulated using D3 branes stretched between NS5branes. Four dimensional theories were discussed in [10,11,12], the most relevant to this review being N = 4 [7], where brane configurations are realized in terms of M theory. In this dissertation we study brane configurations leading to N = 1 supersymmetric theories in five dimensions with gauge group  $SU(N_c)$ and  $N_f$  hyper multiplets, the M theory realization is then described along with its relationship with N = 4 theory in five dimensions.

In section 2 we review the basic properties of five dimensional, N = 1 supersymmetric gauge theories with results established mainly by [1] and [2]. The general prepotential for an N = 1 supersymmetric gauge theory in 5d with  $N_c$  vector multiplets and  $N_f$  hypermultiplets is given[], along with the form it takes when the gauge group is chosen to be  $SU(N_c)$  and the  $N_f$  hypermultiplets transform in the fundamental representation of  $SU(N_c)$ . The conditions on  $N_f$ ,  $N_c$  and  $c_{cl}$  are shown along with the BPS spectrum of the theory. In section 3 we introduce the concept of (p,q) webs in Type IIB string theory [3], starting with how a vertex can be formed from a D5 brane ending on NS5by bending it into a (1, 1) five brane and a general equation is demonstrated for the bending of an NS5 when multiple D5 ending on it . We then move on to (p,q) webs, describing how they can be built to represent  $SU(N_c)$  gauge theories. SU(2) is used as an example to demonstrate how field theoretical results can be obtained from brane configurations. The concept of grid diagrams which are dual to the webs is also explained and how it is used to characterize different theories.

Section 4 starts with a brief description of how Type IIB string theory can be reduced from M theory compactified on a 2-torus [13]. In this context (p,q)branes are M theory five branes wrapped around (p,q) cycles on the torus and brane configurations correspond to an M5 on  $\mathbb{R}^{1,3} \times \Sigma$ , where  $\Sigma$  is a 2 dimensional surface described by a holomorphic curve [5]. We describe how curves for a simple vertex and the SU(2) web are obtained. Grid diagrams are then used to write down curves for pure  $SU(N_c)$  theories. At the end of the section we follow [6] in deriving a general curve for  $SU(N_c)$  theories with  $N_f$  flavors, it is then put into a form where the 4d limit can be taken, and the resulting curve is in analogy with curves given in [8].

## 2 Field Theory content

We start by giving a brief discerption of the field theory, with results mainly established in [1],[2]. We consider N = 1 supersymmetry in five dimensions, which has eight supercharges. The shortest representations are the vector multiplet containing a vector field, a real scalar and fermions, plus a hyper multiplet containing four scalars and fermions.

The Lorentz group is broken to,

$$SO(1,9) \rightarrow SO(1,4) \times SO(3) \simeq SO(1,4) \times SU(2)_R$$
 (2.0.1)

with  $SU(2)_R$  acting as the global R-symmetry group.

Consider a configuration consisting a vector multiplet transforming in the adjoint representation of the gauge group G, and  $N_f$  hypermultiplets transforming in a representation  $R_f$ . The Coulomb branch is parameterized by the Cartan sub algebra of the scalars in the vector multiplet. Low energy dynamics are governed by the prepotential  $\mathcal{F}$  which is restricted to be at most cubic[1], and is given buy [2],

$$\mathcal{F}(\phi) = \frac{1}{2g_0^2} \phi^i \phi_i + \frac{c_{cl}}{2} d_{ijk} \phi^i \phi^j \phi^k + \frac{1}{12} (\sum_{\alpha} |\alpha_i \phi^i|^3 - \sum_f \sum_{w \in R_f} |w_i \phi^i + m_f| \quad (2.0.2)$$

with  $\phi = \phi^i T_i$ ,  $T_i$  are the generators of G in the representation  $R_f$ , and  $d_{ijk} = tr(T_i\{T_j, T_k\})$ . The first sum is over the roots of G while the second is over the weights of  $R_f$ . The bare coupling is  $g_0$  and the effective coupling which is the metric on the Coloumb branch is given by,

$$\left(\frac{1}{g_{eff}^2}\right)_{ij} = \frac{\partial^2 \mathcal{F}(\phi)}{\partial \phi_i \partial \phi} \tag{2.0.3}$$

 $c_{cl}$  is a quantized parameter related to the 5 dimensional Chern-Simons term which characterizes the theory. If we specialize to  $G = SU(N_c)$  with  $N_f$ hypermultiplet in the fundamental representation of G, the moduli space is given by  $\phi = \text{diag}(a_1, a_2 \dots a_{N_c})$  with  $\sum_i a_i = 0$ , modulo the Weyl group which allows us to set  $a_1 \ge a_2 \ge a_3 \dots \ge a_N$ . In this case the prepotential simplifies to[3],

$$\mathcal{F} = \frac{1}{2g_0^2} \sum_{i}^{N_c} a_i^2 + \frac{1}{12} \left( 2\sum_{i>j}^{N_c} |a_i - a_j|^3 + 2c_{cl} \sum_{i}^{N_c} a_i^3 - N_f \sum_{f}^{f} \sum_{i}^{N_c} |a_i + m_f|^3 \right) \quad (2.0.4)$$

In order to have non trivial fixed points for  $N_c > 2$ ,  $N_c, N_f$ , and  $c_{cl}$  have to satisfy the following constraints [2],

$$\frac{1}{2}N_f + c_{cl} \in \mathbb{Z} \tag{2.0.5}$$

$$N_f + 2|c_{cl}| \le 2N_c \tag{2.0.6}$$

A special case is when  $N_c = 2$  where we have two pure gauge theories labeled by their  $\mathbb{Z}$  theta angle,  $d_{ijk} = 0$  and the number of flavors allowed is extended to  $N_f < 7$ . Along with electrically particles, Instanton are present In the BPS spectrum which is charged under the  $U(1)_I$  global symmetries with instanton number I. The central charge is given by,

$$Z = (n_e + I)\phi + m_I$$
 (2.0.7)

masses of BPS states are equal to their central charge, strings are magnetic monopoles with tensions,

$$T_m = (n_m)_i \frac{\partial \mathcal{F}}{\partial \phi_i} \tag{2.0.8}$$

## 3 Type IIB setup

We consider Five brane configurations of Type IIB in the context of  $SL(2,\mathbb{Z})$ , with D5 branes being a (1,0) multiplet, charged under the RR sector 2-form a nd NS5 branes being a (0,1) multiplet, charged under the NS sector 2-form. The D5 branes have world volumes[4],

Brane	World volume	Tension
D5	$(x^0, x^1, x^2, x^3, x^4, x^6)$	$1/g_s l_s^6$
NS5	$(x^0, x^1, x^2, x^3, x^4, x^5)$	$1/g_s^2 l_s^6$

 $g_s$  being the Type IIB string coupling, and  $l_s$  the string length scale. Dimensions  $(x^0, x^1, x^2, x^3, x^4)$  are shared my both NS5 and D5, interesting geometry lies in the  $(x^6, x^5)$  plane, where the branes appear as straight lines. The last three dimensions  $(x^7, x^8, x^8)$  represent the SO(3) R-symmetry and play no role in the Coulomb branch and correspond to deformations in the Higgs branch.

(p,q) branes are formed as bound states of D5 and NS5 branes, their tension being[4],

$$T_{p,q} = |p + \tau q| T_{D5} \tag{3.0.9}$$

#### 3.1 Vertices

A D5 brane ending on an NS5 brane looks like a point charge in one dimension which causes the NS5 brane to bend in to the location,

$$x^{6} = \frac{g_{s}}{2}(\left|x^{5}\right| + x^{5}) \tag{3.1.1}$$

We compose Vertices as our building blocks for the webs to come, by charge conservation they satisfy,

$$\sum_{i} p_i = \sum_{i} q_i = 0 \tag{3.1.2}$$

Following [], a quarter or the supersymmetries are preserved if the branes are constrained to obey a the slope condition,

$$\Delta x_6 + i\Delta x_5 \| p + \tau q \tag{3.1.3}$$

are horizontal while NS5 branes are vertical, the simplest vertex (figure) can be viewed as a D5 ending on an NS5 brane, thus forming a (1,1) brane.



Figure[], vertex extended in the  $(x^6, x^5)$  plane.

#### 3.2 Webs

To generalize (3.1.1) we consider an infinite NS5 brane stretched in  $x^5$  with  $n_L$ D5 branes ending on the left with positions ( in  $x^5$ )  $a_i, i = 1, 2 \dots n_L$  and  $n_R$ branes ending on the right at  $b_i, i = 1, 2 \dots n_R$ , the five brane formed is bent according to [4],

$$x^{6} = \frac{g_{s}}{2} \left( \sum_{i=1}^{n_{L}} |y - a_{i}| - \sum_{i=1}^{n_{R}} |y - b_{i}| + (n_{L} - n_{R}) \right)$$
(3.2.1)

some simple examples are shown below,



 $n_L = 1, n_R = 1$ 



 $n_L = 2, n_R = 1$ 

#### 3.3 Gauge theories

To construct 5 dimensional gauge theories, in this case  $U(N_c)$ , we stretch  $N_c$  parallel D5 branes (in  $x^6$ ) between two NS5 branes. Say we sperate the NS5 branes by  $L_6$  in  $x^6$  and position the D5 branes in  $x^5$  at  $a_i, i = 1, 2 \dots N_c$ , we now have  $N_c - 1$  internal legs between each D5 brane there bending condition can be found by utilizing (3.0.1).

we let the left and right lower external legs to be  $(p_L, q_L)$  and  $(p_R, q_R)$ respectively, by the condition (3.1.2) the left and right internal legs between the *i*'th and i + 1'th D5 branes have charges,

$$(p_L - i, q_L) \tag{3.3.1}$$

$$(p_R + i, q_R) \tag{3.3.2}$$

and the upper external legs have charges,

$$(p_L - N_c, q_L) (3.3.3)$$

$$(p_R + N_c, q_R) \tag{3.3.4}$$

To study gauge theory we need to decouple gravity and massive string modes by considering the limits,

$$L_{max}, g_s, l_s \to 0 \tag{3.3.5}$$

if the gauge coupling is at distance scale L such that,

$$L \gg l_s \gg L_{max} \tag{3.3.6}$$

gravity decouples and massive modes can be integrated out, at low energy this yields a  $U(N_c)$  gauge theory in 1+4 dimensions, with a coulomb branch parameterized by  $a_i, i = 1, 2 \dots N_c$  and coupling,

$$\frac{1}{g_0^2} = \frac{L_6}{g_s l_s^2} \tag{3.3.7}$$

As above, quantum effects bend the branes which restricts the locations of the branes,

$$\sum_{i=1}^{N_c} a_i = 0$$

hence one of the classical positions is fixed and a U(1) factor is frozen, breaking the gauge group to  $SU(N_c)$ .



figure [3], pure SU(2) web.



figure [4], pure SU(3) web.

One can add matter hypermultiplets by having  $N_L$  semi infinite D5 branes to the left of the NS5 branes, and  $N_R$  to the right with,

$$N_L + N_R = N_f \tag{3.3.8}$$

their locations  $m_i, i = 1 \dots f$  in  $x^5$  parameterize the quark masses, the charges of the uppermost left and right five branes are now changed to  $(p_L - N_c + N_L, q_L)$ and  $(p_R + N_c - N_R, q_R)$  respectively.



figure [5], SU(2) with  $N_f = 1$ 



figure [6], SU(3) with  $N_f = 2$ 

An intuitive example that can demonstrate the properties of (p, q) webs is that of pure SU(2) gauge theory (figure). We can immediately read off two field theory parameter, one being the mass of the W boson due to a fundamental string stretched between the two D5 branes,

$$m_W = \Delta x^5 T_s \tag{3.3.9}$$

with  $T_s$  being the string tension and  $\Delta x^5$  the distance between the two D5 branes, which we normalize so that  $m_W = \phi$ , the scalar in our vector multiplet. We also have an instanton due to a D1 string stretched between the two NS5 branes, it has mass,

$$m_I = \Delta x^6 \left| \tau \right| T_s \tag{3.3.10}$$

If we deform the web to a point where the D5 branes ar coincident ( $\phi = 0$ ) we

see that

$$m_I = m_0 = \frac{1}{g_0^2} \tag{3.3.11}$$

At this point (the origin of the moduli space) the W bosons massless and the gauge group is enhanced to SU(2)., however at a generic point of the moduli space ( $\phi \neq 0$ ) the W bosons gain mass,

$$m_W = \phi \tag{3.3.12}$$

and,

$$m_I = \frac{1}{g_0^2} + \phi \tag{3.3.13}$$

the global guage group is now just U(1). The effective coupling is,

$$\left(\frac{1}{g^2}\right)_{eff} = \frac{1}{2}(2m_0 + 4\phi) = m_0 + 2\phi \tag{3.3.14}$$

A monopole corresponding to a D3 brane wrapping the face of the web its tension s given by the area of the face.

$$T_m = \text{area of face} = \phi(\phi + m_0) \tag{3.3.15}$$

We know from field theory results that  $T_m = \frac{\partial \mathcal{F}}{\partial \phi}$  and as,

$$\frac{\partial T_m}{\partial \phi} = (\frac{1}{g^2})_{eff} \tag{3.3.16}$$

the prepotential takes the form,

$$\mathcal{F}(\phi) = \frac{1}{2}m_0\phi^2 + \frac{1}{3}\phi^3 \tag{3.3.17}$$



figure [7], particle states in SU(2).



figure [8], length of dotted red line corresponds to  $1/g_0^2$ 

#### **3.4** Deformations

Deformations of the web play an important role in gauge theory. A **local deformation** is one that does not change the locations of the external legs and hence can be realized by giving VEVs to the scalar fields in the vector multiplet , and hence parameterized by  $\phi$ . The number of local deformations is given by the number of internal faces [3].

$$n_G = \sharp(\text{local deformations}) = \text{rank}(\text{local gauge group}) = \sharp(\text{internal face})$$

$$(3.4.1)$$

and by Euler's formula,

$$n_G = E - V + 1 \tag{3.4.2}$$

where E is the number of internal edges and V is the number of verticies. The metric on the Coulomb branch (effective coupling) is given by the sum of the masses of the edges moved by the deformation, weighted by there displacements squared,



figure [9], a local deformation

Global deformations are ones which do change the asymptotic location of the external legs and there is one deformation associated with each leg. We are only interested in deformations which give new webs so we discard translations in the  $(x^6, x^5)$  plane and an additional deformation constrained by the web. We are left with,

$$n_G = n_X - 3 \tag{3.4.4}$$

with  $n_G$  being the number of global deformations and  $n_X$  the number of external legs. The bare coupling  $\frac{1}{g_0^2}$  can obtained geometrically by deforming the web locally so that the D5 are coincident ( ie.  $\phi = 0$ ),  $\frac{1}{g_0^2}$  is then given by () for the SU(2) case [3],



figure [10], a global deformation

#### 3.5 Grid diagrams

Grids diagrams are an important tool in the classification of five dimensional theories, a thorough description on how to construct grids diagrams is given in [3], here I give a brief description of the concepts and results involved.

A grid diagram is a convex set of points, lines and polygons that lie in a 2-dimensional integer lattice and are labeled by  $(a, b) \in \mathbb{Z}^2$ . Grid diagrams are dual to (p,q) webs in the following way,

- A (p,q) edge in the web is mapped to a vector (-q, p) (line).
- vertices in the web are mapped to polygons.
- Faces are mapped points on the diagram.

By charge conservation at each vertex we have,

$$\sum_{\text{edges}\in\text{vertex}} (p,q) = \sum_{\text{lines}\in\text{polygon}} [-q,p]$$
(3.5.1)

Using a grid diagram one can calculate the dimension of the Coloumb branch by,

$$\dim(\text{Coulomb branch}) = \#(\text{internal points}) \tag{3.5.2}$$

The residual  $SL(2,\mathbb{Z})$  symmetry of the web is carried through by the grid diagram description.



figure[11], a single vertex in the  $(x^6, x^5)$  plane, along with its dual grid diagram in (a, b).

Grid diagrams provide an important insight into the special case of G = SU(2). For an SU(2) we need two parallel D branes which correspond to a column of three points in the grid diagram. The Coulomb branch in one dimensional so we need one internal point and four external points. The left point is fixed by  $SL(2,\mathbb{Z})$  and we are left with 3 possible choices for the point on the left, possible webs are shown below,



figure [12], possible webs for SU(2).

(a) and (b) correspond to theories with theta angles  $\Theta = 0 \pmod{2}$  and  $\Theta = 1 \pmod{2}$  respectively. The web (c) contains paralell external legs which leads

to 6d particles which are charged under global symmetry, if such particles decouple from the low-energy theory (c) will give to a new theory with the same Coulomb branch, but conservatively we assume such theories don't exist.

Pure  $SU(N_c)$  gauge theories can be classified using the concept of grid diagrams, from section (3.3) we need  $N_c$  horizontal D5 branes, which corresponds to a column of  $N_c + 1$  points on the grid. Two additional points are required on both sides of the column, one of which can be fixed by the residual  $SL(2,\mathbb{Z})$ symmetry. There are now  $2N_c+1$  locations one the grid where the point can be placed without breaking convexity, these locations correspond to the possible values of  $c_{cl}$  for  $N_f = 0$ . Charge conjugation  $(c_{cl} \rightarrow -c_{cl})$  can be viewed as a  $\mathbb{Z}_2$  symmetry constructed by an  $SL(2,\mathbb{Z})$  combined with a rotation.



figure[13], possible configurations of SU(3), (\*) point on the left is fixed by  $SL(2,\mathbb{Z})$ , (a)  $c_{cl} = 0$ , (b)  $c_{cl} = N_c$ , (c)  $c_{cl} = -N_c$ 

### 4 Curves

#### 4.1 M theory

To interpret the brane configurations in M theory, we first consider Type IIA string theory in 9+1 dimensions as M theory compactified on a circle in the  $x^{10}$  direction[13]. By comparing brane tensions in Type IIA and M theory, the parameters of the two theories can be mapped to each other by,

$$R_{10} = g_s l_s (4.1.1)$$

$$\frac{R_{10}}{l_p^3} = \frac{1}{l_s^2} \tag{4.1.2}$$

where  $R_{10}$  is the radius of the circle on which  $x^{10}$  is compactified,  $l_p$  being the plank length. In the strong coupling limit  $g_s \to \infty$ ,  $R_{10} \to \infty$  and the theory is described by 10+1 dimensional super gravity. The limit  $g_s, R_{10} \to 0$  takes us to 9+1 dimensional Type IIA string theory.

We further compactify Type IIA on a direction  $x_4$  on circle with radius  $R_A$  by T duality this is equivalent to Type IIB string theory compactified on a radius,

$$R_B = \frac{l_s^2}{R_A} \tag{4.1.3}$$

and string coupling,

$$g'_s = \frac{g_s l_s}{R_A} \tag{4.1.4}$$

This is equivalent to m theory compactified on a 2-torus with sizes  $R_{10}, R_A$ , By (4.1.3) taking the limit  $R_{10}, R_A \to 0$  implies,

$$R_B = \frac{l_p^3}{R_A R_{10}} \to \infty \tag{4.1.5}$$

hence Type IIB is recovered from M theory and the wrapping modes of the M2brane become light and an extra dimension  $x^i$  is created.

In the M theory picture D5 and NS5 branes are interpreted as M5 wrapped around a (1,0) and (0,1) cycles on the torus respectively, (p,q) branes correspond to (p,q) cycles.

#### 4.2 Complex coordinates

 $SL(2,\mathbb{Z})$  acts the complex structure of the 2-torus, hence on the winding numbers (p,q) related to the  $x^4$  and  $x^{10}$  directions. The fact that the orientation of the Type IIB five branes in the  $(x^6, x^5)$  plane means that [5],

$$\Delta x^6 : \Delta x^5 = \Delta x^4 : \Delta x^{10} \tag{4.2.1}$$

This enables us to define complex coordinates,

$$w = x^6 + ix^{10} \tag{4.2.2}$$

$$v = x^4 + ix^5 \tag{4.2.3}$$

as  $x^{10}$  and  $x^6$  are compact it is convenient to define,

$$s = exp(w/R_{10})$$
 (4.2.4)

$$t = exp(-iv/R_A) \tag{4.2.5}$$

Thus in the M theory picture, our brane configuration is described by an M5 brane on  $\mathbb{R}^{1,3} \times \Sigma$ , where  $\Sigma$  is a 2-dimensional surface embedded in  $\mathbb{T}^2 \times \mathbb{R}^2$ 

which is parameterized by s and v. The curve can be written in the form of a holomorphic function F(s,t) such that,

$$F(s,t) = 0 (4.2.6)$$

clearly s = constant and t = constant correspond to NS5 and D5 branes respectively. A simple example would be the curve described by a single vertex, the curve can be computed by considering the asymptotics of s and t [5],

- $s \to 0, x^6 \to -\infty$  so F(s,t) = w 1
- $t \to 0, x^5 \to -\infty$  so F(s,t) = t 1
- $s, t \to \infty, s \sim t$

hence we have [5],

$$F(s,t) = t + s - 1 \tag{4.2.7}$$

(4.2.7) represents a smoothed out vertex in the  $(x^6, x^5)$  plane figure [13], in the 5 dimensional limit  $(R_B \to \infty)$  the vertex and their singularities are retrieved.



figure [13], a simple vertex as a smoothed out M theory curve.

Similarly the curve for SU(2) is given by,

$$F(s,t) = As + t + ABst + Ast^{2} + ts^{2}$$
(4.2.8)

with

$$A \sim 2\exp(L_4/2g_0^2) \quad B \sim 2\exp(m_W L_4/2)$$
 (4.2.9)

#### 4.3 Curves from grids

As was demonstrated in [3], Curves representing a brane configuration can be obtained simply from the grid diagram. Take a (p,q) brane described by the equation,

$$m + (-qx^6 + px^5)T_s = 0 (4.3.1)$$

with m being the transverse position of the brane. Changing variables to (s, t) we have,

$$As^{-q}t^p = 1 (4.3.2)$$

where  $|A| = exp(mR_A^4)$ , this can be rearranged to give,

$$A_1 s^{a_1} t^{b_1} + A_2 s^{a_2} t^{b_2} = 0 (4.3.3)$$

where,

$$(a_1, b_1) - (a_2, b_2) = (-q, p), |A_1/A_2| = exp(mR_A^4)$$
(4.3.4)

So every point (a, b) in the grid diagram can be associated with a monomial  $As^{a}t^{b}$ , for a general grid we have,

$$F(s,t) = \sum_{i \in \text{points on grid}} A_i s^{a_i} t^{b_i}$$

(4.3.5)

coefficients are obtained by exhausting (4.2.11) to a loop of points in the diagram, consider a loop of n points with coefficients  $A_1 \dots A_n$ . We assign  $A_1$ arbitrarily and determine  $A_2$  by,

$$|A_1/A_2| = \exp(m_{1,2}R_A) \tag{4.3.6}$$

with  $m_{12}$  being the transverse position of the brane we cross to get from  $A_1$  to  $A_2$ , similarly,

$$|A_i/A_{i+1}| = \exp(m_{i,i+1}R_A^4) \tag{4.3.7}$$

Tackling the pure  $SU(N_c)$  grid with the same approach, we have a sum of  $N_c + 1$  monomials corresponding to the column of points,

$$(\sum_{i=0}^{N_c} u_i t^i)s$$
 (4.3.8)

the vertical position of the point to the right of the column  $(2, c_{cl})$  characterizes the theory and translate in the curve as the monomial

$$As^2 t^{c_{cl}} \tag{4.3.9}$$

using the residual  $SL(2,\mathbb{Z})$  symmetry the point to the left of the column can be placed at  $(0, c_{cl})$ , putting these results together we have the curve,

$$As^{2}t^{c_{cl}} + (\sum_{i=0}^{N_{c}} u_{i}t^{i})s + Bt^{c_{cl}} = 0$$
(4.3.10)

for constants  $A, u_i, C$ , after some rearrangement and re scaling of s we have,

$$s^2 + P(t)s + 1 = 0 (4.3.11)$$

where

$$P(t) = t^{c_{cl}} (\sum_{i=0}^{N_c} u_i t^i)$$
(4.3.12)

#### 4.4 General $SU(N_c)$

Now we move on to a more general construction given in [6], which will allow us to give restrictions on the possible gauge theories. We study curves that describe SU(N) configurations with  $N_f$  flavors, following []we focus on configurations with two NS5 branes, hence the polynomial will be as most quadratic in t. For convenience we redefine s = exp(-w), the general form of the curve should be

$$A(t)s^{2} + B(t)s + C(t) = 0 (4.4.1)$$

for values of t where C(t) = 0, a solution of the curve is s = 0  $(x^6 \to \infty)$ , hence the zeros of C correspond to semi infinite D5 branes to the right, similarly the zero of A correspond to semi infinite branes to the right and zeros of B correspond the "color" D5 branes.

To normalize the coefficient of  $t^2$  to 1, we move all semi-infinite D5 to the left of the NS5, which corresponds to the transformatio+??n  $s \to \frac{C}{s}$  giving,

$$s^{2} + B(t)s + C(t) = 0 (4.4.2)$$

with

$$B(t) = c_1 t^n \prod_{k=1}^{N_c} (t - a_k)$$
(4.4.3)

and

$$C(t) = c_2 t^m \prod_{k=1}^{N_f} (t - b_k)$$
(4.4.4)

m and n are integers that we will impose conditions on, resealing and taking t to be large  $(x^5 \to \infty)$  we have the leading order polynomial,

$$s^2 + t^{N_c + n}s + t^{N_f + m} = 0 ag{4.4.5}$$

with solution,

$$s = \frac{1}{2} \left( -t^{N_c + n} \pm \sqrt{t^{2(N_c + n)} - 4t^{N_f + m}} \right)$$
(4.4.6)

asymptotically we have the following roots for different  $\,n,m,N_c$  , and  $\,N_f,$ 

- $2(N_c + n) > N_f + m$ ,  $s \sim t^{N_c + n}, t^{-N_c n + N_f + m}$
- $2(N_c + n) < N_f + m$ ,  $s \sim t^{(N_f + m)/2}$
- $2(N_c + n) = N_f + m$ ,  $s \sim t^{N_c + n}$

The case  $2(N_c + n) < N_f + m$  corresponds to NS5 branes crossing, which leads to new massive states being created. This case should be discarded as according to [6] an  $SU(N_c)$  theory with  $N_f > 2N_c$  leads to non trivial fixed points, hence we focus on theories with  $2(N_c + n) \ge N_f + m$ .

Taking the limit  $t \to 0$ , (4.4.2) reduces to,

$$s^2 + t^n s + t^m = 0 (4.4.7)$$

with solution,

$$s = \frac{1}{2}(-t^n \pm \sqrt{t^{2n} - 4t^m})$$

again asymptotically we have,

- $2n > m, s \sim t^{m/2}$
- $2n < m, s \sim t^n, t^{m-n}$
- $2n = m, s \sim t^n$

Taking 2n > m will lead to semi-infinite "flavor" D5 branes crossing, so we only consider  $2n \leq m$ . We still have an  $SL(2,\mathbb{Z})$  transformation which acts as  $t \to tw^l$  taking  $m \to m + 2l$  and  $n \to n + l$ , this allows us to set n = 0, hence,

$$m \ge 0 \tag{4.4.8}$$

which leaves us with the constraint,

$$2N_c \ge N_f + m$$

rearranging we get

$$m \le 2N_c - N_f \tag{4.4.9}$$

under parity  $(t \rightarrow 1/t)$ ,

$$m \to 2N_c - N_f - m$$

hence we are left with  $(2N_c - N_f - 1)/2$  different values for m which correspond to the different values of  $c_{cl}$  found in[2]. The parity operator translates to charge conjugation  $c_{cl} \rightarrow -c_{cl}$ , hence we identify,

$$c_{cl} = N - m - N_f/2 \tag{4.4.10}$$

#### 4.5 Four dimensions

We now turn to reducing our curve to four dimensions, this requires putting (4.4.2) in a form where we can take  $R_A \rightarrow 0$ . Consider,

$$\sin(\frac{v-a_k}{2R_A}) = \frac{1}{2} \left(\exp i(\frac{v-a_k}{2R_A}) - \exp -i(\frac{v-a_k}{2R_A})\right)$$
(4.5.1)

substituting  $t = \exp(-iv/R_A)$  and multiplying through by  $-\exp i(\frac{v}{2R_A})$  we have,

$$-\exp i(\frac{v-a_k}{2R_A})\sin(\frac{v-a_k}{2R_A}) = \frac{1}{2}(t^{1/2} - t^{-1/2}\exp(-i(\frac{a_k}{R_A})))$$
(4.5.2)

multiplying by  $t^{1/2}$  and letting  $\tilde{a}_k = \exp -i(\frac{a_k}{R_A})$  gives,

$$-2t^{1/2}\exp i(\frac{v}{2R_A})\sin(\frac{v-a_k}{2R_A}) = (t-\tilde{a}_k)$$
(4.5.3)

letting  $c_1 = -\exp i(\frac{v}{2R_A})/R_A$  in (4.4.3) we can write B(t) as,

$$B(t) = 2R_A t^{N/2} \prod_{k=1}^{N_c} \sin(\frac{v - a_k}{2R_A})$$
(4.5.4)

similarly for C(t) we have,

$$C(t) = t^{N/2+m} R_A \prod_{k=1}^{N_c} \sin(\frac{v - b_k}{2R_A})$$
(4.5.5)

putting the results together, we can write (4.4.2) as

$$s^{2} + 2sR_{A}t^{N_{c}/2}\prod_{k=1}^{N_{c}}\sin(\frac{v-a_{k}}{2R_{A}}) + t^{N/2+m}R_{A}\prod_{k=1}^{N_{f}}\sin(\frac{v-b_{k}}{2R_{A}}) = 0 \qquad (4.5.6)$$

 $a_k = e^{-i\tilde{a}_k/R_A}$  are know the position of the D5 branes and  $b_k = e^{-i\tilde{b}_k/R_A}$  the positions of the NS5. Taking the 4d limit  $(R_A \to \infty, R_B \to 0)$ , this reduces to,

$$s^{2} + 2s \prod_{k=1}^{N_{c}} (v - a_{k}) + \prod_{k=1}^{N_{f}} (v - b_{k}) = 0$$
(4.5.7)

which is analogous to the curves describing N = 2 SQCD in 4 dimensions[8].

## 5 Conclusion

In this dissertation (p,q) webs were introduced, and how they are used to formulate N = 1 supersymmetric gauge theories. We focus on the case  $G = SU(N_c)$  although concepts could be easily generalized to symplectic and orthogonal groups by the insertion of orientfolds. Grid diagrams are then used to classify different theories, giving the same restrictions that arise in the field theory picture. The M theory description of the brane configurations are then introduced through holomorphic curves which, in the 4d limit, were shown to be analogous to the curves given in[8].

(p,q) webs have provided a very rich venue for studying supersymmetric field theories in five dimensions, it allows us to visualize field theories from a geometrical perspective which paves the way for future work. The interplay between webs, grids and curves is quite enlightening, as it relates three mathematical concepts in the context of field theory.

## References

- Nathan Seiberg, Five Dimensional SUSY Field Theories, "Non-trivial Fixed Points and String Dynamics".hep-th/9608111v2
- [2] Kenneth Intriligator, David R. Morrison, and Nathan Seiberg, "Five-Dimensional Supersymmetric Gauge Theories and Degenerations of CalabiYau Spaces".hep-th/9702198
- [3] Ofer Aharony, Amihay Hanany, and Barak Kol, "Webs of (p,q) 5-branes, Five Dimensional Field Theories and Grid Diagrams".hep-th/9710116v3
- [4] Amit Giveon, "Brane Dynamics and Gauge Theory".hep-th/9802067v2
- [5] Barak Kol, "5d Field Theories and M Theory".hep-th/9705031
- [6] A. Brandhubera, N. Itzhakia, J. Sonnenscheina, S. Theisenb and S. Yankielowizca, "On the M-Theory Approach to (Compactified) 5D Field Theories". hep-th/9709010
- [7] Edward Witten, "SOLUTIONS OF FOUR-DIMENSIONAL FIELD THE-ORIES VIA M THEORY".hep-th/9703166
- [8] Emil J. Martinec and Nicholas P. Warner. "Integrable systems and supersymmetric gauge theory".hep-th/9509161
- [9] Amihay Hanany and Edward Witten"Type IIB Superstrings, BPS Monopoles, And Three-Dimensional Gauge Dynamics".hep-th/9611230

- [10] A. Brandhuber, J. Sonnenschein, S. Theisen and S. Yankielowicz, Brane Configurations and 4D Field Theory Dualities, Nucl. Phys. B502 (1997) 125, hep-th/9704044 c.
- [11] P.C. Argyres, M.R. Plesser, N. Seiberg and E. Witten, New N=2 Superconformal Field Theories in Four Dimensions, Nucl. Phys. B461 (1996) 71, hep-th/9511154.
- [12] C. Ahn and K. Oh, Geometry, D-Branes and N=1 Duality in Four Dimensions I, Phys. Lett. 412B (1997) 274, hep-th/9704061.
- [13] Paul S. Aspinwall, "Some Relationships Between Dualities in String Theory", hep-th/9508154v1.